

Reply to "Comment on 'Scaling of the linear response in simple ageing systems without disorder' "

Malte Henkel¹ and Michel Pleimling²

¹*Laboratoire de Physique des Matériaux (CNRS UMR 7556),*

Université Henri Poincaré Nancy I, B.P. 239, F - 54506 Vandœuvre lès Nancy Cedex, France

²*Institut für Theoretische Physik I, Universität Erlangen-Nürnberg, D - 91058 Erlangen, Germany*

The value of the non-equilibrium exponent a is measured in the two-dimensional (2D) Ising model quenched to below criticality from the dynamical scaling of the zero-field-cooled and the intermediate susceptibility. Our results fully reconfirm the expected value $a = 1/2$ but are inconsistent with the value $a = 1/4$, advocated by Corberi, Lippiello and Zannetti (cond-mat/0506139).

PACS numbers: 05.20.-y, 64.60.Ht, 75.40.Gb

For simple magnets quenched to below their critical point, one generally expects, in the scaling regime where $t, s \gg t_{\text{micro}}$ and also $t-s \gg t_{\text{micro}}$ (t_{micro} is a microscopic reference time), the following scaling behaviour of the linear autoresponse function

$$R(t, s) = \frac{\delta \langle \phi(t) \rangle}{\delta h(s)} \Big|_{h=0} \sim s^{-1-a} f_R(t/s) \quad (1)$$

where $\phi(t)$ is the order parameter at the observation time t and $h(s)$ is the magnetic field at the waiting time s .

For some time, Corberi, Lippiello and Zannetti (CLZ) [1, 2] have advocated a phenomenological formula

$$a = \frac{n}{z} \left(\frac{d - d_L}{d_U - d_L} \right) \quad (2)$$

where $d_{U,L}$ are the upper and lower critical dimensions, respectively and n is the number of components of the order parameter. On the other hand, the commonly accepted physical picture [3] of the ageing (coarsening) process going on after the quench has it that ordered domains form very rapidly and that the ageing comes from the movement of the domain walls between the ordered regimes. If this idea is combined with scaling arguments and if one takes into account that in certain systems (such as the Ising model and referred to as class S) the spatial correlations decay exponentially while in others (such as the spherical model and referred to as class L) the spatial correlations decay algebraically, viz. $C_{\text{eq}}(\vec{r}) \sim |\vec{r}|^{-(d-2+\eta)}$, one obtains [4, 5]

$$a = \begin{cases} 1/z & ; \text{for class S} \\ (d-2+\eta)/z & ; \text{for class L} \end{cases} \quad (3)$$

where z is the dynamical exponent. The result (3) has been reproduced in many studies. However, if eq. (2) of CLZ should turn out to be correct that would also invalidate the simple physical picture of the coarsening process mentioned above. A good practical way to decide between (2) and the prediction (3) appears to be a study of the 2D Ising model with a non-conserved order parameter and quenched to $T < T_c$ from a fully disordered initial state. Here $z = 2$ is known [6] and from eqs. (2) and (3) one has $a = 1/4$ and $a = 1/2$, respectively.

CLZ first arrived at eq. (2) by analyzing numerical data [2] of the field-cooled susceptibility $\chi_{\text{ZFC}}(t, s) = \int_s^t du R(t, u)$. From a straightforward integration of (1), they obtained $\chi_{\text{ZFC}}(t, s) \sim s^{-a}$ and proceeded to extract a . However, it was pointed out that $\chi_{\text{ZFC}}(t, s)$ does not obey a simple scaling but rather there may be a further and dominant contribution coming from the upper limit of integration. For systems of class S, there are well-defined domains with domain walls whose thickness is small with respect to the domain size and one has [5]

$$\chi_{\text{ZFC}}(t, s) = \chi_0 + \chi_1 t^{-A} - s^{-a} f_M(t/s) ; \text{ class S} \quad (4)$$

where $\chi_{0,1}$ are constants and $f_M(x) \geq 0$ is a scaling function. Indeed, it can be shown that $A = z^{-1} - \kappa$ [5], where the exponent $\kappa \geq 0$ describes the time-dependent scaling of the width of the domain walls $w(t) \sim t^\kappa$ [7]. In the 2D Ising model $\kappa = 1/4$ is known [7], hence $A = 1/4$ which explains the early result of CLZ but it becomes clear that the term naïvely expected from the scaling behaviour (1) merely yields a finite-time correction. CLZ did not take into account the condition $t-s \gg t_{\text{micro}}$ for the validity of the scaling form (1) in their analysis [2] and hence have missed the leading time-dependent term in (4).

On the other hand, for systems of class L, we argued previously that because of the long-range correlations, the effective width of the domains should be the same as their linear size which leads to $A = 0$ [5]. Since the clusters have no 'inside', the term analogous to $\chi_0 = 0$ in eq. (4) is absent, but it's rôle is taken over by the constant χ_1 . CLZ [1] point out correctly that for both classes S and L the constant $\lim_{t \rightarrow \infty} \chi_{\text{ZFC}}(t, s)$ (with $x = t/s$ fixed) and which equals χ_0 for class S and $\chi_0 + \chi_1$ for class L can be related to the equilibrium magnetization through the static fluctuation-dissipation theorem. In summary one has, with a given by (3)

$$\chi_{\text{ZFC}}(t, s) = \begin{cases} \frac{1}{T} (1 - m_{\text{eq}}^2) + \chi_1 t^{-A} - s^{-a} f_M(t/s) & ; \text{for class S} \\ \frac{1}{T} (1 - m_{\text{eq}}^2) - s^{-a} f_M(t/s) & ; \text{for class L} \end{cases} \quad (5)$$

In their comment, CLZ [1] now argue that the exponents A and a were really not distinct and that one rather




FIG. 1: (a) Scaling plot of the reduced field-cooled susceptibility $\chi_{\text{ZFC}}(t, s) - \chi_0$ against t/s in the 2D Ising model quenched to $T = 1.5$, for waiting times $s = 25, 50, 100, 200, 800, 1600$ (from bottom to top). (b) Comparison of the data with a fit $\chi_{\text{ZFC}}(t, s) - \chi_0 = a_0 s^{-1/4} + a_1 s^{-1/2}$ (black lines) and a fit $\chi_{\text{ZFC}}(t, s) - \chi_0 = a_0 s^{-1/4}$ (grey lines), for several values of t/s . For clarity the data for $t/s = 2$ and for $t/s = 3$ have been shifted upwards by a factor 1.4 and 1.2, respectively.

had $a = A$. From now on, we concentrate on systems of class S and furthermore on the 2D Ising model quenched below T_c . From numerical simulations, CLZ reproduce once more that $\chi_{\text{ZFC}}(t, s) - \chi_0 \sim s^{-1/4}$, as generally expected. But they do not consider explicitly the corrections to that leading scaling behaviour and merely claim, but do not prove, the absence of the scaling corrections of order s^{-a} , which are expected from eq. (5).

Indeed, the scaling behaviour of χ_{ZFC} is not as simple as claimed by CLZ. We show this in figure 1a, with data obtained from the standard heat-bath method. It can be seen that the collapse in this scaling plot is far from complete which strongly indicates the presence of sizeable finite-time corrections. The origin of these becomes clear in figure 1b where we compare our data to the asymptotic form $\chi_{\text{ZFC}}(t, s) - \chi_0 = a_0 s^{-1/4} + a_1 s^{-1/2}$, for several values of t/s . While taking into account both terms does reproduce the data well, we included for comparison also the fits where $a_1 = 0$ was fixed by hand, as suggested by CLZ. It is clear that the data, which display a pronounced curvature, can only be fitted if $a_1 < 0$, as expected from (5). Indeed, the best fits yield $a_0 = 0.68$ and $a_1 = -0.35$ for $t/s = 2$, $a_0 = 0.62$ and $a_1 = -0.19$ for $t/s = 3$ and $a_0 = 0.54$ and $a_1 = -0.06$ for $t/s = 5$ in excellent quantitative agreement with (5). We conclude that there is no indication in favour of an absence of the last term in eq. (5), hence $a > A$ which provides clear counterevidence against the proposal of CLZ.

We had already given earlier an analysis of the scaling of the thermoremanent magnetization (where terms of order s^{-A} do not occur). By taking into account also the

leading finite-time correction we found full compatibility with $a = 1/2$ but inconsistency with $a = 1/4$ [4].

We may also show that indeed $a = 1/2$ in the 2D Ising model through the so-called intermediate integrated response [5, 8]

$$\chi_{\text{Int}}(t, s) = \int_{s/2}^s du R(t, u) \sim s^{-a} f_{\text{Int}}(t/s) \quad (6)$$

which has the advantage that the leading term $\sim t^{-A}$ which dominates over the scaling term in χ_{ZFC} is absent. In figure 2 we examine the scaling of $\chi_{\text{Int}}(t, s)$ and try to achieve a collapse for $a = 1/2$ and $a = 1/4$. While there is a nice collapse for $a = 1/2$ already for relatively small values of the waiting time (unless t/s is too close to unity), the data fail to collapse for $a = 1/4$. Again, this is fully consistent with $a = 1/2$ but excludes $a = 1/4$.




FIG. 2: Scaling plot of the intermediate susceptibility $\chi_{\text{Int}}(t, s)$ for (a) the assumed value $a = 1/2$ (b) the assumed value $a = 1/4$. Error bars are smaller than the symbol sizes.

Turning to systems of class L, CLZ nicely reconfirmed the expected scaling forms eqs. (3,5). However, there is nothing in their test with contradicts the simple scaling we used in [5] to obtain $A = 0$.

In conclusion, having reexamined the scaling of some integrated susceptibilities, we have shown that the scaling of $\chi_{\text{ZFC}}(t, s)$ does indeed contain at least two important contributions, see eq. (5), which is against the proposal of CLZ. We stress that a simple demonstration of scaling of χ_{ZFC} is not enough to be able to reliably know which of the exponents A or a is measured. It is more safe to study a quantity such as the intermediate susceptibility, which does not suffer from this difficulty. Applied to the Ising model quenched to $T < T_c$, our results fully confirm eq. (3) but disagree with eq. (2).

This work was supported by CINES Montpellier (projet pmn2095). MP was supported by the Deutsche Forschungsgemeinschaft through grant no. PL 323/2.

[1] F. Corberi, E. Lippiello, and M. Zannetti, preceeding comment and references therein.

[2] F. Corberi, E. Lippiello, and M. Zannetti, Eur. Phys. J.

- B24**, 359 (2001); Phys. Rev. **E68**, 046131 (2003).
- [3] see the reviews by A.J. Bray and J.P. Bouchaud in M.E. Cates and M.R. Evans (eds), *Soft and Fragile Matter*, IOP Press (Bristol 2000).
- [4] M. Henkel, M. Paessens and M. Pleimling, Europhys. Lett. **62**, 664 (2003).
- [5] M. Henkel, M. Paessens and M. Pleimling, Phys. Rev. **E69**, 056109 (2004).
- [6] A.J. Bray, Adv. Phys. **43**, 357 (1994).
- [7] D.B. Abraham and P.J. Upton, Phys. Rev. **B39**, 736 (1989).
- [8] M. Pleimling and A. Gambassi, Phys. Rev. **B71**, 180401(R) (2005).